

Neurosurgical Methods **Luis Rafael Moscote-Salazar**

Neurosurgery is a medical specialty that deals with the prevention, diagnosis and treatment of disorders affecting nervous system including brain, peripheral nerves, spinal cord, and extra-cranial cerebrovascular system. **Insights in Neurosurgery** is an Open access, peer reviewed international journal that publishes scientific articles related to all aspects of neurology, focusing mainly on neuro surgery. The current volume no. 1, issue 2 has published three research articles, three case reports and a commentary.

Jha et al. in their research article concluded that patients suffering from dorsal Pott's disease with antero-lateral or anterior compression with significant neurological deficit needs Unilateral transpedicular decompression with bed rest for three months for recovery [1]. Oppido and Cattani in their studies on various methods of neuroendoscopy concluded that Tu laser is extremely handy, safe and a reliable dissecting method for neuroendoscopy [2].

Konig and Spetzger in their research article compared the effectiveness of cervical total disc replacement (TDR) and anterior cervical discectomy and fusion (ACDF). The study found no significant difference between TDR group and ACDF group patients in the clinical outcomes even after 2 years of surgery [3]. Acar et al. successfully treated a 57 year old women suffering from Parkinson's disease by using Deep brain stimulation technique, which alleviates motor symptoms and complications [4].

Sbeih et al. [5] reported the case of Cerebral Toxoplasmosis

Neurosurgeon-Critical Care, RED LATINO,
Latin American Trauma & Intensive Neuro-
Care Organization, Bogota, Colombia

Corresponding author:

Luis Rafael Moscote-Salazar

✉ mineurocirujano@aol.com

Neurosurgeon-Critical Care, RED LATINO, Latin
American Trauma & Intensive Neuro-Care
Organization, Bogota, Colombia.

Tel: +57301283538

Citation: Moscote-Salazar LR. Neurosurgical
Methods. Neurosurg. 2016, 1:2.

in a 40 year old married male Syrian with HIV and Tominga et al. [6] reported a Transient Bilateral Oculomotor Nerve Palsy (TOP) in a 67 year old male which is associated with ruptured anterior communicating artery aneurysm. Cuoco and Davy briefly discussed about the ethical considerations in Human head transplantation [7].

REFERENCES

- 1 Jha DK, Singh R, Pant I (2016) Transpedicular surgical decompression of dorsal spinal tuberculosis (pott 's disease) with vertebral collapse without fixation. *Neurosurg 1*: 1-8.
- 2 Oppido PA, Cattani F (2016) The utility of thulium laser in neuroendoscopy. *Neurosurg 1*: 1-4.
- 3 König SA, Spetzger U (2016) Clinical outcome of anterior cervical discectomy and fusion versus total disc replacement – A meta-analysis of 2532 cases. *Neurosurg 1*: 1-6.
- 4 Acar G, Göçmen S, Kocaoğlu M, Civlan S, Zisakis A, et al. (2016) Bilateral pallidotomy by using deep brain stimulation electrode in Parkinson's disease. *Neurosurg 1*: 1-4.
- 5 Sbeih I, Bilbisi M, Darwazeh R, Al-gahoom T, Abu-Farsakh H (2016) Cervical spinal cord toxoplasmosis in a patient with aids: Case report and review of the literature. *Neurosurg 1*: 1-7.
- 6 Tominaga T, Aihara Y, Kawamata T, Arai K (2016) Transient bilateral oculomotor nerve palsy (top) associated with ruptured anterior communicating artery aneurysm: A case report. *Neurosurg 1*: 1-5.
- 7 Cuoco JA, Davy JR (2016) Operation Frankenstein: Ethical reflections of human head transplantation. *Neurosurg 1*: 1-4.